

My patient reached out after testing positive for COVID-19. How do I help them?

1 Confirm that your patient:

- Has been diagnosed with mild to moderate COVID-19 with positive results of direct SARS-CoV-2 viral testing
- Is within 10 days of symptom onset
- Is an adult or pediatric patient (age 12-17 years and weighing at least 40 kg)

2 Assess if patient is at high risk for progressing to severe COVID-19 and/or hospitalization.

- The following medical conditions or other factors may place adults and pediatric patients (age 12-17 years and weighing at least 40 kg) at higher risk for progression to severe COVID-19:
 - > Older age (for example age ≥ 65 years of age)
 - > Obesity or being overweight (for example, adults with BMI > 25 kg/m², or if age 12-17, have BMI ≥ 85 th percentile for their age and gender based on [CDC growth charts](#))
 - > Pregnancy
 - > Chronic kidney disease
 - > Diabetes
 - > Immunosuppressive disease or immunosuppressive treatment
 - > Cardiovascular disease (including congenital heart disease) or hypertension
 - > Chronic lung diseases (for example, chronic obstructive pulmonary disease, asthma [moderate to severe], interstitial lung disease, cystic fibrosis, and pulmonary hypertension)
 - > Sickle cell disease
 - > Neurodevelopmental disorders (for example, cerebral palsy) or other conditions that confer medical complexity (for example, genetic or metabolic syndromes and severe congenital anomalies)
 - > Having a medical-related technological dependence (for example, tracheostomy, gastrostomy, or positive pressure ventilation [not related to COVID-19])
- Other medical conditions or factors (for example, race or ethnicity) may also place individual patients at high risk for progression to severe COVID-19 and is not limited to the medical conditions or factors listed above. For additional information on medical conditions and factors associated with increased risk for progression to severe COVID-19, see the [CDC website](#). Healthcare providers should consider the benefit-risk for an individual patient.

3 Identify an infusion center near your patient.

- Call the Lilly COVID Hotline at 1-855-545-5921. Explain to one of our dedicated medical professionals that you are a healthcare provider and that one of your patients is COVID-positive and meets the criteria for use of monoclonal antibodies. Ask the agent to provide you the name and number of the infusion site nearest the patient.

4 Write the order.

- Contact the infusion site to understand the preferred process at that site for prescribing monoclonal antibody treatments.

5 Provide your patient with resources found on [LillyAntibody.com](#) (click on FAQs & Resources) so that he or she arrives at the appointment feeling prepared. For additional information, healthcare providers and patients can also call the Lilly COVID Hotline at 1-855-545-5921.